

Contoocook Valley School District

FOOD/INSECT & EMERGENCY ALLERGY CARE PLAN and MEDICATION AUTHORIZATION

STUDENT INFORMATION	Student Name	DOB:
	Home/Cell Phone	Grade
	Known Life-Threatening Allergies:	History of Asthma? <input type="checkbox"/> No <input type="checkbox"/> Yes (Asthma may indicate an increased risk of severe reaction)
	Diagnosis of Oral Allergy Syndrome? <input type="checkbox"/> No <input type="checkbox"/> Yes	History of SEVERE Anaphylactic Reaction? <input type="checkbox"/> No <input type="checkbox"/> Yes,
	Please list allergens:	If checked YES, give epi-pen immediately if allergen was <i>likely</i> eaten, at onset of <i>any</i> symptoms, and follow the protocol below

TREATMENT PLAN	<p>ANY ONE OF THESE SEVERE SYMPTOMS OF ANAPHYLAXIS AFTER SUSPECTED OR KNOWN INGESTION:</p> <ul style="list-style-type: none"> ➤ Difficulty breathing or swallowing ➤ Dizzy, faint, confused, pale or blue, hypotension/weak pulse <p style="text-align: center;">OR</p> <p>ANY COMBINATION OF SYMPTOMS FROM DIFFERENT BODY AREAS:</p> <p>AIRWAY: Short of breath, chest tightness, wheeze, repetitive cough, profuse runny nose</p> <p>THROAT: Tight, hoarse, trouble breathing/swallowing, drooling</p> <p>MOUTH: Swollen lips or tongue</p> <p>SKIN: Hives, Itchy rashes, swelling (e.g., eyes, lips)</p> <p>GUT: Nausea, Vomiting, diarrhea, cramp-like pain</p>	<p>FOLLOW THIS PROTOCOL:</p> <ol style="list-style-type: none"> 1. INJECT EPINEPHRINE IMMEDIATELY! 2. Call 911 3. Raise feet above the head, remain lying down & continue monitoring 4. Give additional medications as ordered <ul style="list-style-type: none"> - Antihistamine - Bronchodilator/Albuterol if has asthma 5. Notify Parent/Guardian 6. Notify Prescribing Provider / PCP 7. When indicated, assist student to rise slowly.
	<p>ORAL ALLERGY SYNDROME (IF DIAGNOSIS CONFIRMED ABOVE):</p> <p>MOUTH: Itchy mouth, lips, tongue and/or throat</p> <p>SKIN: Itching just around mouth</p>	<ol style="list-style-type: none"> 1. GIVE ANTIHISTAMINE (swish, gargle, & swallow) 2. Monitor student as indicated; notify healthcare provider & parent as indicated 3. If progresses to symptoms of anaphylaxis, USE EPINEPHRINE (as stated above)

➤ THE SEVERITY OF SYMPTOMS CAN QUICKLY CHANGE. ALL SYMPTOMS OF ANAPHYLAXIS CAN POTENTIALLY PROGRESS TO A LIFE THREATENING SITUATION!!

DOSEAGE OF MEDICATIONS	Epinephrine	<input type="checkbox"/> Epi Auto-injector, Jr (0.15mg) inject intramuscularly <input type="checkbox"/> Epi Auto-injector (0.3mg) inject intramuscularly ➤ A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur.	
	Antihistamine	<input type="checkbox"/> Benadryl/Diphenhydramine Dose: Route: PO Frequency:	<input type="checkbox"/> Other Dose: Route:
	Medication shall be administered during school year:	TO	<p>NOTE: IF NURSE IS NOT AVAILABLE, THE EPINEPHRINE AUTO INJECTOR MAY BE GIVEN BY DESIGNATED SCHOOL PERSONNEL FOR ANY ANAPHYLAXIS SYMPTOMS</p>

TO BE COMPLETED BY PARENT AND AUTHORIZED HEALTHCARE PROVIDER

AUTHORIZATION	<p>Prescriber's Signature: _____ Prescriber's Authorization to Self-Administer</p> <p><i>Confirms student is capable to safely and properly administer medication</i> <input type="checkbox"/> Yes <input type="checkbox"/> No</p>	<p>Date: _____</p> <p>PRESCRIBER'S PRINTED NAME OR STAMP AND SIGNATURE:</p>
	<p>Parent: I hereby request that the above ordered medication be administered by school personnel and consent to communications between the school nurse and the prescriber that are necessary to ensure safe administration of this medication. This protocol will be in effect until the end of the current or extended school year. This medication will be destroyed if not picked up within one week following termination of the order or the end of the school year. Whichever comes first, unless the student will be attending an extended school year (ESY) program. A new protocol will be needed for the next school year. I have received, reviewed and understand the above information.</p>	
	<p>Parent's Signature: _____ Parent's Authorization to Self-Administer</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No</p>	<p>Date: _____</p>

EMERGENCY CARE PLAN FOR STUDENT

NAME: _____ GRADE/SCHOOL: _____

Insert Picture if available

SYMPTOMS OF ANAPHYLAXIS:

- Chest tightness, shortness of breath, cough, wheezing, profuse runny nose
- Dizzy, faint, pale, blue, confused
- Tightness and/or itching in throat, difficulty swallowing, hoarseness, drooling
- Swelling of lips, tongue, throat
- Itchy mouth, itchy skin, hives
- Hives, itching (anywhere), swelling (eg face, eyes)
- Nausea, vomiting, diarrhea, cramp-like pain

IF ALLERGEN LIKELY EATEN (OR STUDENT STUNG), FOLLOW THIS EPINEPHRINE PROTOCOL AT THE ONSET OF ANY OF THE ABOVE SYMPTOMS:

1. Administer Epi Auto-Injector: **circle one:** (0.15mg 0.3mg)
2. Have someone call 911 for ambulance, don't hang up, and stay with student
3. Administer Benadryl: **circle one** 12.5mg 25mg 37.5mg 50mg other _____
4. Have student lie down with feet above level of head until EMS arrives
5. Notify school and parent/guardian as soon as possible

EPI AUTO-INJECTOR DIRECTIONS:



For EPIPEN and EPIPEN JR.:

1. Pull off blue activation cap.
2. Hold orange tip near outer thigh (always apply to thigh). Okay to inject through clothing.
3. Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10; remove and massage 10 sec. Auto-Injector should then be removed and take to Emergency Room.



For Auvi-Q:

1. Follow verbal instructions.
2. Pull off red safety guard. Pull firmly to remove.
3. Place black end against middle of outer thigh (through clothing if needed.) Then press firmly and hold in place for 5 seconds.



EMERGENCY CONTACTS

1. Name: _____
Relation: _____
Phone: _____

2. Name: _____
Relation: _____
Phone: _____

EMERGENCY/PHYSICIAN CONTACTS

1. Name: _____
Relation: _____
Phone: _____

2. Name: _____
Relation: _____
Phone: _____

Parent

Student (if applicable)

School Nurse